


Luxembourg

Total population: 524,000 Total deaths: 3,600 Income group: High Life expectancy at birth: Total:82 Males:80 Females:84





Age-Standardized Cancer Mortality Trends





Cancer Incidence





Adult Risk Factors

| | Males | Females | Total |
|--|-------|---------|-------|
| Current tobacco smoking (2011) | 22.3% | 25% | 23.7% |
| Total alcohol per capita consumption, in litres of pure alcohol (2010) | 16.8 | 7.2 | 11.9 |
| Physical inactivity (2010) | 28.2% | 31.7% | 30.0% |
| Obesity (2014) | 28.3% | 21.3% | 24.8% |
| Household solid fuel use (2012) | - | - | 0.0% |

| Cancer Plans, Monitoring and Surveillance | | |
|---|---|--|
| Has an operational cancer policy/strategy/action plan | No | |
| Has a cancer registry | Yes | |
| Scope | Population-based | |
| Coverage | National | |
| Last year of data | NR | |
| Cancer Primary Prevention Policies | | |
| Tobacco control | | |
| Has an operational policy, strategy or action plan to reduce the burden of tobacco use | Yes | |
| Smoke-free legislation | Data not categorized | |
| Tobacco dependence treatment | NRT and/or some cessation services (at least one of which is cost-covered | |
| Warning labels | Medium size warnings missing some appropriate characteristics OR large warnings missing many appropriate characteristics | |
| Bans on advertising, promotion and sponsorship | Ban on national television, radio and print media as well as on some but no all other forms of direct and/or indirect advertising | |
| Tobacco taxes | 51–75% of retail price is tax | |
| Overweight and obesity prevention and control | | |
| Has an operational policy, strategy or action plan for reducing overweight/obesity | Yes | |
| Physical inactivity prevention and control | | |
| Has an operational policy, strategy or action plan to reduce physical inactivity and/or promote physical activity | Yes | |
| Harmful use of alcohol prevention and control | | |
| Has an operational policy, strategy or action plan to reduce the harmful use of alcohol | No | |
| National immunization | | |
| Human Papillomavirus vaccination schedule | 12-18 years | |
| Hepatitis B vaccination schedule | | |
| Hepatitis B vaccination coverage, infants | 94% | |
| Cancer Screening and Early Detection Cervical cancer | | |
| Cervical cytology (PAP) | Generally available at the public primary health care level | |
| Acetic acid visualization (VIA) | Generally available at the public primary health care level | |
| Breast cancer | asilotally arailable at the position primary results date to to | |
| Breast palpation / clinical breast exam (CBE) | Generally available at the public primary health care level | |
| Mammogram | Generally available at the public primary health care level | |
| Colorectal cancer | 1 | |
| Faecal occult blood test or faecal immunological test | Generally available at the public primary health care level | |
| Bowel cancer screening by exam or colonoscopy | Generally available at the public primary health care level | |
| Cancer Treatment and Palliative Care | | |
| Radiotherapy | Generally available in the public health system | |
| Total high energy teletherapy units / million inhabitants | 3.8 | |
| Number of radiotherapy centres | 1 | |
| Number of radiation oncologists | 4 | |
| Chemotherapy (medicines not specified) Oral morphine (formulation not specified) | Generally available in the public health system Generally available in the public health system | |
| Non-methadone morphine equivalent consumption per cancer death (mg) | 61,680.7 | |
| Community/home care for people with advanced stage capear and other NCDs | Generally available | |

Generally available

Community/home care for people with advanced stage cancer and other NCDs