


Niger

Total population: 17,157,000 Total deaths: 180,000 Income group: Low Life expectancy at birth: Total:59 Males:59 Females:59





Age-Standardized Cancer Mortality Trends*





Cancer Incidence





Adult Risk Factors

| | Males | Females | Total |
|--|-------|---------|-------|
| Current tobacco smoking (2011) | 8.9% | <1% | 4.8% |
| Total alcohol per capita consumption, in litres of pure alcohol (2010) | 0.5 | 0.1 | 0.3 |
| Physical inactivity (2010) | 21.2% | 24.8% | 23.0% |
| Obesity (2014) | 1.7% | 5.7% | 3.7% |
| Household solid fuel use (2012) | - | - | 94.0% |

| Cancer Plans, Monitoring and Surveillance | | | |
|---|---|--|--|
| Has an operational cancer policy/strategy/action plan | No | | |
| Has a cancer registry | Yes | | |
| Scope | Population-based | | |
| Coverage | Subnational | | |
| Last year of data | 2012 | | |
| Cancer Primary Prevention Policies | | | |
| Tobacco control | | | |
| | No | | |
| Has an operational policy, strategy or action plan to reduce the burden of tobacco use | No | | |
| Smoke-free legislation | Three to five public places completely smoke-free | | |
| Tobacco dependence treatment | NRT and/or some cessation services (neither cost-covered) | | |
| Warning labels | Large warnings with all appropriate characteristics** | | |
| Bans on advertising, promotion and sponsorship | Ban on all forms of direct and indirect advertising** | | |
| Tobacco taxes | 26–50% of retail price is tax | | |
| Overweight and obesity prevention and control | | | |
| Has an operational policy, strategy or action plan for reducing overweight/obesity | No | | |
| Physical inactivity prevention and control | | | |
| Has an operational policy, strategy or action plan to reduce physical inactivity and/or promote physical activity | No | | |
| Harmful use of alcohol prevention and control | | | |
| Has an operational policy, strategy or action plan to reduce the harmful use of alcohol | No | | |
| National immunization | | | |
| Human Papillomavirus vaccination schedule | 11 years, +3 months, +7 months (subnational) | | |
| Hepatitis B vaccination schedule | | | |
| Hepatitis B vaccination coverage, infants | 70% | | |
| Cancer Screening and Early Detection | | | |
| Cervical cancer | | | |
| Cervical cytology (PAP) | Not generally available at the public primary health care level | | |
| Acetic acid visualization (VIA) | Not generally available at the public primary health care level | | |
| Breast cancer | | | |
| Breast palpation / clinical breast exam (CBE) | Generally available at the public primary health care level | | |
| Mammogram | Not generally available at the public primary health care level | | |
| Colorectal cancer | | | |
| Faecal occult blood test or faecal immunological test | Not generally available at the public primary health care level | | |
| Bowel cancer screening by exam or colonoscopy | Not generally available at the public primary health care level | | |
| Cancer Treatment and Palliative Care | | | |
| Radiotherapy | Not generally available in the public health system | | |
| Total high energy teletherapy units / million inhabitants | 0.0 | | |
| Number of radiotherapy centres | | | |
| Number of radiation oncologists | | | |
| Chemotherapy (medicines not specified) | Not generally available in the public health system | | |
| Oral morphine (formulation not specified) | Not generally available in the public health system | | |

^{*} No mortality data available. Figures are based on national incidence estimates and modelled survival.
** Indicates highest possible level of achievement

Non-methadone morphine equivalent consumption per cancer death (mg) Community/home care for people with advanced stage cancer and other NCDs

Not generally available

^{... =} No data available