

PAKISTAN


CERVICAL CANCER PROFILE

TOTAL POPULATION,
FEMALE (2019): **105 100 000**

TOTAL DEATHS,
FEMALE (2019): **741 000**


Morbidity and Mortality

Crude cervical cancer incidence per 100 000 women (2020):	4.7
Age-standardized cervical cancer incidence per 100 000 women (2020):	6.1
Cumulative risk of cervical cancer, ages 0-74 (2020):	0.7%
Cervical cancer deaths (2019):	4 000
Cervical cancer mortality-to-incidence ratio (2020):	0.64
Population-based cancer registry exists (2021):	No



Primary Prevention

HPV vaccination programme coverage among girls (2020)



HPV vaccination is not included in the national vaccination schedule

HPV vaccination programme (2020):

HPV included in national vaccination programme:	No
Scale of vaccination programme:	-
Year of introduction:	-
Primary target cohort:	-


Related risk factors:

Tobacco use prevalence, women aged 15+ years (2020):	7%
Condom use at last high-risk sex (-):	ND
HIV incidence per 1000, women aged 15+ years (2020):	0.07

Secondary Prevention

National screening programme for cervical cancer exists (2021):	No
Primary screening test used (2021):	-
Target age range of programme (2021):	-
Programme/guidelines exist to strengthen early detection of first symptoms at primary health care level (2021):	No
Clearly defined referral system exists from primary care to secondary and tertiary care (2021):	No

Screening for cervical cancer (2019)



Fewer than 1 in 10 women have been screened for cervical cancer in the last 5 years

Treatment and Supportive Care

National guidelines on cervical cancer management exist (2021):	No
---	-----------

Number of radiotherapy units per 10 000 cancer patients (2021):	4
Number of brachytherapy units per 10 000 cancer patients (2021):	1

Cancer diagnosis and treatment services generally available (2021):

Cancer centre or cancer department at tertiary level:	Yes
Pathology services (laboratories):	Yes
Cancer surgery:	Yes
Chemotherapy:	Yes
Radiotherapy:	No

Number of medical staff (per 10 000 cancer patients):

Radiation oncologists (2019):	2
Medical physicists (2019):	ND
Surgeons (2012):	129
Radiologists (2019):	57
Nuclear medicine physicians (2019):	14

Palliative care for patients with NCDs in the public health system generally available (2021):

In primary health care facilities:	No
In community or home-based care:	No

Reported annual opioid consumption - excluding methadone - in oral morphine equivalence per capita (2017): **<1mg**

WHO Cervical Cancer Elimination Strategy Targets for 2030

90% of girls fully vaccinated with the HPV vaccine by the age of 15

70% of women are screened with a high-performance test by 35 years of age and again by 45 years of age

90% of women identified with cervical disease receive treatment

ND = data not available

* **Programme coverage:** % of national target population (among 9-14-year-old girls)

† **Coverage by age 15:** % of population turning 15 that have been vaccinated against HPV at any time between ages 9 to 14

[See Explanatory Notes for indicator descriptions.](#)