SERBIA

CERVICAL CANCER PROFILE


TOTAL POPULATION. FEMALE (2019): 4 475 000

TOTAL DEATHS.

FEMALE (2019): 57 200


Morbidity and Mortality





Primary Prevention

HPV vaccination programme coverage among girls (2020)



HPV vaccination is not included in the national vaccination schedule

HPV vaccination programme (2020):

HPV included in national vaccination programme:	No
Scale of vaccination programme:	-
Year of introduction:	-
Primary target cohort:	-


Related risk factors:

Tobacco use prevalence, women aged 15+ years (2020)**:	36%
Condom use at last high-risk sex (-):	ND
HIV incidence per 1000, women aged 15+ years (2020):	<0.01

Secondary Prevention

National screening programme	
for cervical cancer exists (2021):	Yes
Primary screening test used (2021):	cytology
Target age range of programme (2021):	25-64 years
Programme/guidelines exist to strengthen early detection	
of first symptoms at primary health care level (2021):	Yes
Clearly defined referral system exists from primary care	
to secondary and tertiary care (2021):	Yes

Screening for cervical cancer (2019)



6 in 10 women have been screened for cervical cancer in the last 5 years

Treatment and Supportive Care

National guidelines on cervical cancer management	
exist (2021):	Yes

Number of radiotherapy units per 10 000 cancer patients (2021): 6 Number of brachytherapy units per 10 000 cancer patients (2021):

Cancer diagnosis and treatment services generally available (2021):

Cancer centre or cancer department at tertiary level:	Yes
Pathology services (laboratories):	Yes
Cancer surgery:	Yes
Chemotherapy:	Yes
Radiotherapy:	Yes

Number of medical staff (per 10 000 cancer patients):

Radiation oncologists (2019):	ND
Medical physicists (2019):	10
Surgeons (-):	ND
Radiologists (2019):	ND
Nuclear medicine physicians (2019):	13

Palliative care for patients with NCDs in the public health system generally available (2021):

In primary health care facilities:	Yes
In community or home-based care:	Yes

Reported annual opioid consumption excluding methadone - in oral morphine equivalence per capita (2017):

WHO Cervical Cancer Elimination Strategy Targets for 2030

90% of girls fully vaccinated with the HPV vaccine by the age of 15

70% of women are screened with a high-performance test by 35 years of age and again by 45 years of age

90% of women identified with cervical disease receive treatment

ND = data not available

* Programme coverage: % of national target population (among 9-14-year-old girls) † Coverage by age 15: % of population turning 15 that have been vaccinated against HPV at any time between ages 9 to 14

**Data refer to tobacco smoking only, in absence of sufficient data on all tobacco use